

Ageratum conyzoides L.: A review on its phytochemical and pharmacological profile

Anjoo Kamboj, Ajay Kumar Saluja¹

Guru Gobind Singh College of Pharmacy, Yamuna Nagar - 135 001, Haryana, India, ¹A.R. College of Pharmacy, Vallabh Vidhyanagar - 388 120, Gujarat, India

Ageratum conyzoides L. (AC) is an annual herbaceous plant with a long history of traditional medicinal uses in many countries in the world, especially in the tropical and subtropical regions. The weed has been known since ancient times for its curative properties and has been utilized for treatment of various ailments, such as burns and wounds, for antimicrobial properties, for many infectious conditions and bacterial infections, arthrosis, headaches and dyspnea, pneumonia, analgesic, anti-inflammatory, antiasthmatic, antispasmodic and haemostatic effects, stomach ailments, gynaecological diseases, leprosy and other skin diseases. A wide range of chemical compounds including alkaloids, coumarins, flavonoids, chromenes, benzofurans, sterols and terpenoids have been isolated from this species. Extracts and metabolites from this plant have been found to possess pharmacological and insecticidal activities. This contribution provides a comprehensive review of its ethnomedical uses, chemical constituents and the pharmacological profile as a medicinal plant. Particular attention has been given to analgesic and anti-inflammatory, antibacterial and wound healing properties, radioprotective, antifungal, anthelmintic and nematocidal, insecticidal effects presented in this review such that the potential use of this plant either in pharmaceuticals or as an agricultural resource can be evaluated.

Key words: *Ageratum conyzoides*, alkaloids, benzofurans, bioactivity, chromenes, flavonoids, terpenoids

INTRODUCTION

Ageratum is derived from the Greek words 'a geras', meaning non-aging, referring to the longevity of the whole plant. *Conyzoides*, on the other hand, is derived from 'konyz' the Greek name of *Inula helenium*, which the plant resembles.^[1] The plant belongs to the family Asteraceae tribe Eupatorieae. This family is well marked in their characteristics and cannot be confused with any other. The genus *Ageratum* consists of approximately 30 species but only a few species have been phytochemically investigated.^[2]

It is a tropical plant that is very common in West Africa, Australia, some parts of Asia and South America. It is an erect, annual, branched, slender, hairy and aromatic herb, which grows to approximately 1 m in height. The stems and leaves are covered with fine white hairs, the leaves are stalked, ovate, 4-10 cm long and 1-5 cm wide, with tip and base somewhat pointed and with round-toothed margins long. The flowers are purple to white, less than 6 mm across and arranged in close terminal inflorescences. The fruit is black and are easily dispersed while the seeds are photoblastic and often lost within 12 months.^[3] The plant grows commonly in waste and on ruined sites. It has a peculiar odour likened in Australia to that of a male goat and hence its name 'goat weed' or 'billy goat weed'.

The essential oil obtained from plant has been reported to have a powerful nauseating odour^[4] and found to be poisonous to rabbits due to the presence of HCN and coumarin.^[5] The herb is not eaten by humans except for medicinal purposes, but in some cultures, it is a delicacy for domestic guinea-pigs, horses, cattle,^[6] and also used to feed fish.^[7] Presence of Pyrrolidine alkaloids are known to be hepatotoxic and to cause lung cancer and other ailments^[8] but of biological interest due to their association with Leptidoptera.^[9]

ETHNOPHARMACOLOGY

Ageratum conyzoides (AC) has been used in various parts of Africa, Asia, and South America for curing various diseases. Githen, in his review^[10], listed the uses of the plant as purgative, febrifuge, for ophthalmia, colic, treatment of ulcers, and wound dressing. The antienteralgic and the antipyretic properties of the plant were also indicated in a review on 'Medicinal plants from Senegal'.^[11] In some African countries, the plant has been popular use for skin diseases, wound healing, mental and infectious diseases, headaches and dyspnea,^[12,13] and used in traditional medicine for its anti-asthmatic, antispasmodic and haemostatic effects,^[14] uterine troubles, pneumonia by rubbing them on the chest of the patient.^[15] In Cameroon, it is a local remedy for craw-craw.^[5]

In India, it is used in the treatment of leprosy and as an oil lotion for purulent ophthalmia.^[16] The decoction or infusion

For correspondence: Anjoo Kamboj, GGS College of Pharmacy, Yamuna Nagar - 135 001, Haryana, India.

E-mail: anjoo73_kamboj@indiatimes.com

Received: 02-12-2007; **Accepted:** 28-01-2008

of the herb is given in stomach ailments such as diarrhoea, dysentery, intestinal colic, flatulence, rheumatism, fever^[17] and to relieve pain associated with navel in children.^[15] In Brazil, medicinal tea plants are used as anti-inflammatory, analgesic and anti-diarrhoeic^[18] and for gynaecological diseases (in Vietnam).^[19] Other folk remedies include anti-itch, sleeping sickness, mouthwash for toothache, antitusive, vermifuge, tonic and killing lice (flowers).^[20] The leaves are used for application on cuts, sores;^[21-25] as anti-inflammatory agent, haemostatic;^[26-30] as an insecticide;^[31] in headache;^[20,32] in boils;^[33] skin diseases;^[34] ringworm infection;^[35] in typhoid, as an antidote to snake poison;^[36,37] in malarial fever, as antitetanus; for uterine problems;^[38,39] prolaps of anus;^[40] for swollen piles;^[41] in throat infection, painful gums; on abscess for early suppuration; in wound healing and leucorrhoea.^[42-45] The root is used as an antilithic^[21,26,38] and in infant diarrhoea.^[46] It has been reported to have nematocidal activity and have potential use in controlling pests.^[2,47] It is the only plant used in HIV/AIDS disease.^[48]

Ageratum conyzoides also has a number of magical and superstitious attributes, e.g. in Ivory Coast, it has protective fetish properties for followers of Snake-Sect against snakebite. In Congo, the leaf sap on the hands of card players is believed to improve their luck. If sap is spread on the accused in a trial and is then pricked with a needle, pain will be felt only if guilty.^[2]

PHYTOCHEMISTRY

The oil content varies randomly from 0.11 to 0.58% for leaves and from 0.03 to 0.18% for the roots depending on times of the year.^[49] From water distillation of the fresh flowers, the oil content was found to be 0.2%.^[4] The yield of oil from the petroleum ether extract of the seed was 26%.^[50]

Mono and Sesquiterpenes

The GC-MS analysis of the essential oil showed that it is a complex mixture of 213 compounds of which 43-51 constituents have been reported. The constituents identified include 20 monoterpenes 6.4% [13 monoterpene hydrocarbons (5.0%), 7 oxygenated monoterpenoids (0.08-1.4%)], 20 sesquiterpenes 5.1% [16 sesquiterpene hydrocarbons (4.3%), four oxygenated sesquiterpenoids (0.8%)] and three phenylpropanoids and benzenoids (2.33%). The most abundant monoterpenes include sabinene, β -pinene 1.6%, β -phellandrene, 1,8-cineole and limonene 2.9%, terpinen-4-ol 0.6%, α -terpineol, linalool 0.5%, α -pinene 6.6%, eugenol 4.4% and methyleugenol 1.8%. Sesquiterpenes hydrocarbons (26.48% of the oil) contain β -caryophyllene (19.79%), caryophyllene epoxide (0.5%), β -sesquiphellandrene (1.2-1.99%), δ -cadinene (4.3%) and τ -cadinene (1.44%) as the major components.^[51-53] The major

sesquiterpenes are β -caryophyllene 1.9%, 10.5% from the oil obtained from Cameroon^[51,54] and 14-17% in Pakistani oil.^[55] The oxygenated sesquiterpene hydrocarbons (69.29% of oil) comprised ageratochromene (32.90%), 6-methoxyquinoline-1-oxide (20.77%), β -caryophyllene oxide (7.29%) and β -sinensal (5.82%) as the major components.^[51,52] Ocimene is found in trace amount in Nigerian plant and 5.3% of the oil from Indian plant.^[53]

Chromene, Chromone, Benzofuran, and Coumarin

The most common components of the essential oil are 7-methoxy-2,2-dimethylchromene (precocene I) ranging from 30 (Vietnamese oil) to 93% (Congo oil), 6,7-dimethoxy derivative, ageratochromene (precocene II) ranges from 0.7 to 55%^[49,56-58] and ageratochromene dimer.^[16-59] Other related compounds include enecalinal, 6-vinyl-7-methoxy-2,2-dimethyl chromene, dihydroencecalinal, dihydrodemethoxyencecalinal, demethoxyencecalinal, demethylencecalinal and 2-(1'-oxo-2'-methylpropyl)-2-methyl-6,7-dimethoxychromene.^[60,61] In addition, seven other chromene derivatives isolated from hexane extract of the aerial part of the plant are 2,2-dimethylchromene-7-O- β -glucopyranoside,^[61,62] 6-acetyl-2,2'-dimethyl-3,4-dihydrochromene,^[51] 6-(1-methoxyethyl)-7-methoxy-2,2-dimethylchromene, 6-(1-hydroxyethyl)-7-methoxy-2,2-dimethylchromene, 6-(1-ethoxyethyl)-7-methoxy-2,2-dimethyl chromene, 6-angeloyloxy-7-methoxy-2,2-dimethylchromene and an inseparable mixture of enecanescins.^[63] Benzofuran derivatives, 2-(2'-methylethyl)-5,6-di methoxybenzofuran,^[60,61] 14-hydroxy-2H β , 3-dihydroeuparine,^[62] 6,7,6',7'-tetramethoxy-2,2,2',2'-tetramethyl-3'(4')-dehydro-3'-4S-bi chroman^[16,59] as well as chromone derivatives, 3-(2'-methyl propyl)-methyl-6,8-dimethoxychrom-4-one and 2-(2'-methylprop-2'-enyl)-2-methyl-6,7-dimethoxychroman-4-one have also been reported from the plant.^[60,61] The essential oil from Brazil has been reported to yield 1.24% of coumarin.^[64] The ethanolic extract of the plant was found to be devoid of tannins.^[65]

Flavonoids

Ageratum conyzoides is very rich in polyoxygenated flavonoids. A total of 21 polyoxygenated flavonoids have been reported. This includes 14 polymethoxylated flavones that are tricin derivatives, 3',4',5'-oxygenated flavones, which include ageconyflavone A, ageconyflavone B and ageconyflavone C, other flavonoids include 5'-methoxynobiletin, linderoflavone B, 5,6,7,3',4',5'-hexamethoxyflavone, 5,6,8,3',4',5'-hexamethoxyflavone, eupalestin, nobiletin, 5,6,7,5'-tetramethoxy-3',4'-methylenedioxyflavone, sinensetin, 5,6,7,8,3',5'-hexamethoxyflavone, 5,6,7,8,3'-pentamethoxy-4'-hydroxyflavone and 5,6,7,8,3',5'-hexamethoxy-4'-hydroxyflavone. Conyzorigun originally believed to be a phenoxy chromone was found to be identical with eupalestin.^[164] The polyhydroxyflavones

include scutellarein-5,6,7,4'-tetrahydroxyflavone, quercetin, quercetin-3-rhamnopyranoside, kaempferol, kaempferol-3-rhamnopyranoside and kaempferol-3,7-diglucopyranoside. A novel isoflavone glycoside, 5,7,2',4'-tetrahydroxy-6,3'-di-(3,3-dimethylallyl)-isoflavone 5-O- α -L-rhamnopyranosyl-(1 \rightarrow 4)- α -L-rhamnopyranoside isolated from the stems.^[57,63,66-74]

Triterpene and Sterols

The triterpenes isolated from plant are friedelin, and the major sterols are sterols- β -sitosterol (26.7%) and stigmasterol (59.9%). Other minor sterols isolated are brassicasterol (0.3%) and dihydrobrassicasterol (2.7%), spinasterol (5.2%), dihydrospinasterol (5.7%). The leaves also contain a rare sterol, stigmast-7-en-3 β -ol.^[75-79]

Alkaloids and Miscellaneous Compounds

Lycopsamine and echinatine, two isomeric pyrrolizidine alkaloids, are the only alkaloids isolated from this plant.^[78] Other compounds isolated from plant include (+)-sesamin,^[63] aurantiamide acetate,^[79] fumaric acid, caffeic acid,^[73] phytol, HCN and hydrocarbons, from nC₂₇-H₅₆ to nC₃₂-H₆₆.^[9] (Z) 12-(6-methyl)-heptadecenoic acid.^[80] The seed oil was found to contain oleic acid (25.52%),^[81] palmitic, stearic, linoleic, linolenic and hexadecenoic acids.^[50] The amino acid composition in the leaves and flowers are cystine, leucine, histidine, arginine, serine, alanine, tyrosine, valine, phenylalanine and threonine, glycine (trace).^[82] The amino acid composition of the pollen was found to 0.5-4.0% of the total dry weight. They are amino-n-butyric acid, aspartic acid, cystine, isoleucine, methionine, proline, tryptophan, tyrosine, ornithine, glutamic acid and glycine.^[83] The protein content found in the leaves is 4.69% and in flowers 9.37%. Leaves and flowers contain in addition fructose, ribose and glucose and vitamin B while vitamin A was reported to be present in the flowers.^[84,85]

BIOACTIVITY

Ageratum conyzoides is a plant used in traditional medicine for mental and infection diseases, dyspnea, enteralgia and fever. The plant was reported to have significant antifungal and insecticidal properties. The biological activities of the plant are treated under two subheadings: (i) pharmacological properties and (ii) insecticidal and other biological properties.

PHARMACOLOGICAL PROPERTIES

Crude Extract

The crude extract of the whole plant has been reported to be superior to vaseline gauze as a wound dressing material.^[70] It has been found to have neuromuscular blocking activity in isolated rat's phrenic nerve-diaphragm and

also caused greater fall in diastolic pressure compared with that of systolic pressure in anaesthetized rats. It has calcium blocking activity similar to that of verapamil.^[86] The leaf extract has been used in the treatment of chronic pain in osteoarthrotic patients.^[87] It has antimicrobial and anticonvulsant activities.^[14,88] Aqueous extract of the leaves has been reported to prevent coagulation of the whole blood while causing precipitation of some blood materials and the bleeding time was also decreased.^[89]

Cardiovascular Activities

The water soluble portion of the plant potentiated pentobarbitone-induced hypnosis in mice in a dose of 1 g/kg. It produced hypotensive response in dog, which was not blocked by propranolol, anthesisan or atropine. The extract revealed negative inotropic and chronotropic effects in doses of 10-40 mg on frog's perfused heart, which was not blocked by atropine. It showed a direct vasodilatory response in rats, and its smooth muscle activity similar to papaverine. The extract had no analgesic and anticonvulsant activity.^[90] The essential oil in a concentration of 0.1 ml/ml caused contraction of the isolated frog rectus abdominis within 15 s after administration and showed a graded response up to a dose of 0.3 ml. The plant (leaves and root) was devoid of *in vivo* cholinesterase activity.^[91]

The aqueous extract of leaves is reported to show haemostatic activity. The activity is observed probably due to vasoconstriction and the formation of an artificial clot that tend to produce a mechanical plug to arrest bleeding from small blood vessels. The plant extract exhibited muscle-relaxant activity on isolated rabbit ileum and cardio-depressant activity on isolated heart.^[89,92] The leaves extract changes the electrocardiogram, atrial impulse velocity and coronary vessel resistance on isolated guinea-pig heart.^[93]

Analgesic and Anti-inflammatory Activities

Clinical trials in patients with arthrosis have been conducted with the aqueous extract of the whole plant. Result shows analgesic effect in 66% of patients and improvement in articulation mobility in 24% without side effect.^[94] In another study, the analgesic activity of the leaf extract was detected by hot plate method.^[95] The extract decreased spontaneous motor activity and caused a fall in rectal temperature. *In vitro* receptor radio ligand assay was carried out on the extract to demonstrate its selectivity to a single receptor implicated in the mediation of pain. Result showed more than 50% inhibition in the bradykinin (BK II) assay, but the activity was lost after PVP treatment suggesting that phenolic compounds could be responsible for the initial bioactivity.^[96] The extracts, however, did not produce positive effects in the neurokinin (NKI) and calcitonin gene-related peptide (CGRP) assays.

The antinoceptive properties were analyzed in the wistar rat by using Martin *et al.*, method slightly modified. The rectal temperature was sampled and analgesic activity evaluated by the hot plate method and by the writhing syndrome induced by acetic acid. The results obtained demonstrate that the doses of 50 and 100 mg/kg are selected for experiment it causes: a precocious ataxia, sedation and a slight ptosis for 6 h after injection.^[18,95]

Aqueous extract of an association of AC, *Cymbopogon citrates* and *Lippia multiflora* produced a significant reduction in mouse of writhings induced by acetic acid and an increase of pain threshold in the hot plate test in mice. Presence of saponins and flavonoids supports the observed activities and suggests that this association of three plants could be used as traditional improved preparation.^[97]

The water soluble fraction of the plant extract produces peripheral analgesic activity and an anti-inflammatory action, which seems to occur in leucocyte-dependent inflammatory events.^[98,99] Silva *et al.*, investigating the effect on smooth muscles using isolated rat uterus and intestine smooth muscles, concluded that the fraction possesses substances, which provoke direct relaxing effect on smooth muscles and inhibit contraction induced by several agonists possibly by blocking the entry of calcium and/or inhibiting cAMP phosphodiesterase.^[100] These pharmacological characteristics could explain the popular use of AC to alleviate abdominal and menstrual pains. The aqueous extract of leaves was evaluated for their cytoprotective activity against ethanol-induced gastric lesions in rats. Microscopically, pretreated rats with aqueous extract or cimetidine showed significantly marked inhibition of gastric lesions and marked reduction of submucosal oedema compared to control group.^[101]

The ethanolic root extract of plant in a dose of 100 and 300 mg/kg significantly reduced the carrageenin-induced hind paw oedema in rats with no acute toxic effect in mice.^[102] The hydroalcoholic extract of leaves was studied for its anti-inflammatory effect on subacute (cotton pellet-induced granuloma) and chronic (formaldehyde-induced arthritis) models of inflammation in rats and evaluated by biochemical and haematological analysis of rats blood samples. The results confirm the anti-inflammatory properties with no apparent hepatotoxicity.^[103] The ethanolic extract also showed significant gastroprotective activity that could be mediated by its antioxidant activity, Ca²⁺ channel blocking and antiserotonergic properties.^[104] The ethanolic extract of roots possesses anti-inflammatory and analgesic properties.^[102]

Antibacterial and Wound Healing Properties

The antibacterial properties of methanolic extract were

studied against 11 wound isolates *S. aureus* (four strains), *E. coli* (two strains), *P. aeruginosa* (one strain), *Proteus* spp. (three strains) and *Shigella* spp. (one strain) using the well diffusion method. Wound healing properties were determined using the excision wound model. More than 90% wound healing was recorded in the extract, whereas 72% healing was observed in the distilled water-treated group.^[105,106] The leaf extract was found to be active against *C. falcatum* and *R. solani*^[107] and showed strong toxicity against the fungi causing ringworm, *E. floccosum*, *T. mentagrophytes* and *M. gypseum*, the inhibition of the mycelia being 80.28, 78.43 and 68.24%, respectively.^[108] The extract of leaves, however, had no effect on the conidial germination of the fungus *D. oryzae*.^[109] In a study, it was found that wounds treated with aqueous leaves extract in combination with honey and with solcosery ointment significantly accelerate wound healing process and the rates of wounds sterility compared to wounds treated with honey alone.^[110,111]

The Nigerian traditional soft soaps prepared using varied locally sourced raw materials have been evaluated antimicrobial activities using standard pharmacopoeia protocols and an *in vitro* agar diffusion bioassay method. The incorporated medicinal plants used did not show any significant effect on the antimicrobial activities exhibited by the various soaps against the bacterial and fungal test organisms.^[112]

Antibacterial activity of aqueous extract were tested against three Gram-positive bacteria and seven Gram-negative bacteria and evaluated by the filter paper disc diffusion method. Results showed a significant control of the growth of *A. viscolactis*, *K. aerogenes*, *B. cereus* and *S. pyogenes*.^[113,114,161] In an investigation for *in vitro* anti-methicillin-resistant *Staphylococcus aureus* (MRSA) activity found that MIC range of 55.4-71.0 mcg/ml were recorded for ethanol and water extracts of AC. The concentrations were too high to be considered active and AC were found to be ineffective *in vitro* in the study; therefore, suggest the immediate stoppage of their traditional use against MRSA-associated diseases in Lagos, Nigeria.^[115]

Radioprotective Activity

The effect of alcoholic extract of the plant on the alteration of radiation-induced mortality in mice exposed to 10 Gy of gamma radiation was studied. The mice were treated with 75 mg/kg (optimum dose) AC extract before exposure to 6, 7, 8, 9, 10 and 11 Gy of gamma radiation. The AC extract treatment effectively protected mice against the gastrointestinal as well as bone marrow-related death. In an *in vitro* study, AC extract was found to scavenge 1,1-diphenyl-2-picrylhydrazyl (DPPH) radicals in a concentration-dependent manner, indicating that the

radioprotection may be due to the scavenging of reactive oxygen species induced by ionizing radiation.^[116]

Reproductive Problems

Many herbal remedies are traditionally used as contraceptives, abortifacients, emmenagogues or oxytocics. Forty-two plants are used for reproductive problems of men and women. AC is used to treat the prostate problems, unspecified female complaints and venereal disease. The AC extract inhibited uterine contractions induced by 5-hydroxy tryptamine suggesting that the extract exhibited specific antiserotonergic activity on isolated uterus but had no effect on uterine contractions induced by acetylcholine. The results support the popular use of the plant as a spasmolytic.^[117,162]

Essential Oil

Essential oil of AC has been tested for anti-inflammatory, analgesic and antipyretic activities in mice and rats. The oil was found to have a significant anti-inflammatory cotton pellet granuloma activity and the antipyretic Brewer's yeast injection effect, whereas the analgesic tail-flick and writhing test activity with no gastric toxicity.^[118]

Antibacterial and Antifungal Activities

The essential oil obtained from the plant showed antibacterial activity against *V. cholerae*, *S. shigae*, *S. pyogenes*, *C. diphtheriae* and *S. typhi*,^[119,120] whereas it was devoid of any activity against the potato pathogenic bacteria viz., *B. polymyxa*, *E. carotovora* var. *atroseptica*, *E. carotovora*, *P. putida*, *P. solanacearum* and *X. vesicatoria*.^[113] In other study, oil exhibit antibacterial and antifungal activities against 22 bacteria, including Gram-positive cocci and rods and Gram-negative rods and 12 fungi (three yeast-like and nine filamentous).^[121] The total inhibition of growth was recorded against the four fungi, *C. albicans* SP-14, *C. neoformans* SP-16, *S. rolfisii* SP-5 and *T. mentagrophytes* SP-12. Rao^[122] reported that the oil inhibited the growth of five bacteria as well as 10 fungi species and that the major component of the oil demethoxyageratochromene^[1] was effective against two of the fungi, *P. chrysogenum* and *P. javanicum*.^[51] The oil provided 100% inhibition of the mycelial growth and germination of spores of *Didymella bryoniae*.^[123] The volatile oil extracted from the leaves possesses molluscicidal property.^[51,124]

Industries have researched on the use of essential oils as an alternative for the treatment of some microbiologic pathologies. *A. flavus* is a fungus present in food and feeds and can produce aflatoxins under some environmental conditions. The effect of the essential oil and crude ethanolic extract on *A. flavus* growth was carried out by the disc diffusion method. The essential oil of AC was able to inhibit *A. flavus* growth, but crude ethanolic extract was not effective against the tested fungal in the concentration

used. In another study, the essential oil of the plant showed antifungal activity against *H. turcicum*, *H. oryzae*, *C. capsici*, *P. setariae* and *F. moniliforme*.^[114,125] At a concentration of 2000ppm, the oil exhibit strong antifungal activity against *E. floccosum*, *M. canis* and *T. mentagrophytes*.^[126]

Anthelmintic and Nematicidal Activity

The essential oil of the plant exhibited anthelmintic activity against *T. solium* and *P. posthuma*.^[127] The extract of the stem and leaves exhibited nematicidal activity against second stage juveniles of *Meloidogyne incognita* (kofoid and white) chitwood.^[128,129]

Metabolites

Pharmacological activities of the most significant metabolites from this plant, responsible for the medicinal properties, have not been identified. There are however, a wide spectrum of pharmacological activities of the classes of compounds obtained from this plant. For example, simple chromenes and chromans, especially the 6-amino and 6-acetamido derivatives, have been reported to have anti-depressant, analgesic and antipyretic properties. Some of them possess activity against flat worms of the order of trematodes.^[130] Other simple 2,2-dimethyl chromene derivatives, such as 6-(1-hydroxyethyl)-7,8-dimethoxy-2,2-dimethylchromene and 6-hydroxy-7,8-dimethoxy-2,2-dimethyl chromene, have been shown to have antimicrobial activities.^[131]

The sterols, especially stigmasterol, have been shown to exert significant anti-inflammatory activity.^[132] The flavonoids possess a wide range of biological activities that include effects on CVS, diuresis, spasmolytic, antiviral and anti-inflammatory properties.^[15,19] The free radical scavenging and anti-cancer activities of the flavonoids are well known. Two polymethoxyflavones are found to have potent activity for inducing differentiation of human promyelocytic leukemia cells (HL-60). Although the biological activities of flavonoids isolated from AC have not been investigated, the polymethoxyflavones are important candidates for cancer-protective action.^[133]

Insecticidal and Other Biological Properties

Insecticidal activity

Ageratum conyzoides has bioactivity that may have agricultural use. The crude plant extract also showed insecticidal and pesticidal activities against various types of insects and pests. The major components of the oil, namely the precocenes, have been reported to have antijuvencidal hormonal activity.^[163] The oil exerted acute toxicity on adults of cowpea weevil, *Callosobruchus maculatus* F., upon fumigation. Application of oil dressing on cowpea seed exhibited insecticidal activity against weevil. Significant oviposition deterrence and complete inhibition of emergence of adult insects F1 offspring from oil-treated

beans were evident at 2.5-10 µl/9.5 g beans with no adverse physiological effect. Precocene I was found to be four times as active as the oil.^[134]

Assays conducted in India showed high nymphal mortality 91% of the oil to the Nymphs of *S. gregaria*.^[61] Calle *et al.* showed that the hexane extract of the whole plant showed activity against *M. domestica* larvae.^[135] Methanolic extract from fresh leaves (250 and 500 ppm) also produced deficiency of juvenile hormone in the fourth instar of *C. partellus*, a sorghum pest.^[136] Antijuvénile hormonal activity of Precocenes I and II have been demonstrated on a variety of insects, which include *S. oryzae*, *T. japonica*, *L. chinensis* and *D. flavidus*.^[137] The results from these assays include precocious metamorphosis of the larvae, production of sterile, moribund and dwarfish adults. The two chromenes have been reported to act synergistically and they survived metabolism for at least 12 days.^[137] Preliminary study on the mode of action of precocene II on *M. domestica* L. and *L. caesar* L. has been carried out.^[138] While the precocenes have been seen as fourth-generation insecticides, the drawback is that they have been shown to cause hepatotoxicity in rats.^[139-141] This is an important factor bearing in mind the human health hazard in field applications of precocenes as large-scale insecticidal agents. Some workers demonstrated that the toxicity was due to a highly reactive precocene-3,4-epoxide, a metabolite produced in insect species from cytochrome P-450.^[139,140] Others, like Darvas *et al.* and Casas *et al.*^[142,143] reported that the 3,4 double bond played no significant role in the toxicity but that the oxidative dealkylation process at C7 position, as tocopherol-like antioxidants, might be responsible for the cytotoxicity.

The insecticidal activities of hexane and ethanol extract were evaluated against adults of *R. dominica*. Only the hexane extract showed insecticidal activity. 5,6,7,8,3',4',5'-Hepta methoxyflavone showed low activity against *D. hyalinata* and *R. dominica* and was not toxic to *M. domestica* or *P. americana*. In contrast, coumarin showed insecticidal activity against all four insect and pest species tested.^[144] The extract of the plant was found to exhibit juvenilizing effect on *D. cingulatus* nymphs.^[65]

The alcoholic extract of the plant possessed insecticidal activity against *M. domestica* and *T. castaneum*.^[145] The petroleum ether and acetone extract of the plant showed juvenile hormone activity against *C. quinquefasciatus*, *A. aegypti* and *A. stephensi*.^[146,147] The methanolic extract of the plant was found to suppress the population of the malaria vector *A. stephensi* in higher dosage, whereas in lower dosage it was found to induce several developmental defects and ultimately decrease the growth index to a considerable extent.^[146] In another study, higher concentrations of the crude extract suppressed the vector population of

C. quinquefasciatus, whereas the lower concentrations were found to induce developmental defects ultimately decreasing the growth index of the treated second and fourth instar larvae.^[63,96,127,148]

The petroleum ether extract of the leaves, flowers and buds diluted in benzene and mixed with green gram seeds proved very repulsive, a potent oviposition inhibitor and a safe protectant against the infestation of pulse beetle *C. chinensis*.^[149] The crude extract of the plant showed insecticidal activity against nymphs of mustard aphid *L. erysimi*.^[150]

Allelopathy and Allelopathic Potential

Both the volatile oil and the aqueous extract of AC have been shown to have allelopathic effects on a number of cultivated crops. These include radish, mungbean and ryegrass.^[151] The saturated aqueous solution of the isolated and purified precocenes I and II have been reported to have significant inhibitory effect on the seedling growth of radish, tomato and ryegrass.^[152] The allelopathic potential of the aqueous extract from different organs of AC and from its different development stages, especially from different habitats, was different.^[153,160]

The inhibitory effects of AC volatiles on peanut, redroot amaranth, cucumber and ryegrass increased when plants were grown under nutrient-deficient conditions or in competition with *B. pilosa*; however, there was no difference with physical damage or 2,4-D treatment. Phytoinhibitory effects decreased under fungal infection and aphid feeding. Volatiles from AC plants infected with *E. cichoracearum* or exposed to *A. gossypii* feeding inhibited or killed fungi and insects.^[154]

Phytotoxic Action

Ageratum conyzoides showed strong inhibition on *R. sativus*, *L. germination* and growth in a bioassay. The leaves exhibited a greater suppression than the stem and root. Application of AC leaves at 2 t/ha in a paddy field 2 days after transplanting caused about 75% paddy weed reduction and increased yield by 14% compared with an herbicide treatment. The greater number of growth inhibitors (phenolic compounds) found in the leaves might result in the stronger inhibitory activity than the stem and root. AC might be a natural herbicide for weed control in paddy fields to reduce the dependence on synthetic herbicides.^[155]

Toxicological Studies

The pollination season of the plant causing allergy has been found to be between November and January in Delhi. In another study, *Ageratum* (species not mentioned) pollen were reported to be cause of nasobronchial allergy in 5 out of 50 patients.^[156]

CONCLUDING REMARKS

The genus *Ageratum* is a rapidly spreading plant and is presently a major problem for environmentalists, ecologists, farmers and animal scientists. A number of studies have been carried out on its control as a weed.^[157,158] Of all the species, AC has been widely studied. It is believed that detailed information as presented in this review on its phytochemistry and various biological properties of the extracts and the constituents might provide incentive for proper evaluation of the use of the plant in medicine and in agriculture. Some small companies in Brazil are reported to be using AC as raw material for phytochemicals. Test conducted in mice and rats for anti-inflammatory, analgesic, antipyretic, antibacterial, antifungal, anti-ulcer, radioprotective activities have shown significant results without adverse side effects. Similarly, clinical trials with arthrosis patients conducted with the aqueous extract of the whole plant did not show any side effect.^[87] The major constituents of the essential oil have been shown to produce precocious metamorphosis in insect larvae as well as sterility, moribund and dwarfishness in adult insects. Further work, however, still needs to be carried out on the toxicity of the plant and, especially on the precocenes, which have been shown in few cases to cause hepatotoxicity in rats.^[139,140] The flavonoids possess wide range of biological activities like its effects on CVS, diuresis, antiviral, spasmolytic, anti-inflammatory properties of the flavonoids isolated from the plant need to be studied. Precocenes and coumarins have been seen as fourth-generation insecticides, also need to be studied. Germane to the pest control studies is the report that the plant extract induces morphogenetic abnormalities in the formation of mosquito larvae *C. quinquefasciatus*, *A. aegypti* and *A. stephensi*.^[146,159] Further studies in this area could serve as a means of controlling the Anopheles mosquitoes and thus prevent human and economic losses caused by malaria.

REFERENCES

- Kissmann G, Groth D. Plantas infestantes e nocivas. Sao Paulo:Basf Brasileira; 1993.
- Burkill HM. The Useful Plants of West Tropical Africa, vol. 1. Kew: Royal Botanic Gardens; 1985.
- Marks MK, Nwachuku AC. Seed bank characteristics on a group of tropical weeds. Weed Res 1986;26:151-7.
- Sood VK. Chemical examination of the flower oil of *Ageratum conyzoides* L. Flav Ind 1973;4:77.
- Abbiw DK. Useful plants of Ghana, Intermediate Tech. Publication. London: Royal Botanic Gardens, Kew; 1990. p. 207.
- Dalziel JM. The useful plants of West Africa. London: Crown Agents; 1937.
- Menut C, Lamaty G, Amvam PH. Aromatic plants of tropical central Africa part X: Chemical composition of essential oil of *Ageratum houstonianum*. Flav Frag J 1993;8:1.
- Couett CE, Crews C, Hanley BA. Analysis, separation and bioassay of pyrrolizidine alkaloids from comfrey. Nat Toxins 1996;4:1605-10.
- Vera R. Chemical composition of essential oil of *Ageratum conyzoides* L. from Reunion. Flav Fragr J 1993;8:257.
- Githens TS. Drug plants of Africa. Afr Handbooks 1948;8:59.
- Kerharo J, Adam JG. La pharmacopée senegalaise traditionnelle: Plantes. Medicinales et toxiques vol. 1, Paris: Vigot; 1974. p. 101.
- Adjanohoun EJ, Ahyi AM, Ake-Assi L. Médecine traditionnelle et pharmacopée, 1988.
- Durodola JI. Antibacterial property of crude extracts from a herbal wound healing remedy *Ageratum conyzoides* L. Planta Medica 1977;32:388-90.
- Kokwaro JO. Medicinal plants of East Africa. Nairobi: East African Literature Bureau; 1976. p. 58.
- Okunade AL. Ph.D Thesis, Nigeria: University of Ibadan; 1981. p. 84.
- Katsuri TR, Thomas M, Abraham EM. Essential oil of *Ageratum conyzoides*: Isolation and structure of two new constituents. Indian J Chem 1973;11:91-5.
- Chopra RN, Nayar SL, Chopra IC. Glossary of Indian Medicinal Plants. New Delhi: NISCIR; 2002. p. 9.
- Yamamoto LA, Soldera JC, Emim JA, Godinho RO, Souccar C, Lapa AJ. Pharmacological Screening of *Ageratum conyzoides* L. Mem Inst Oswaldo Cruz 1991;86:145-7.
- Sharma PD, Sharma OP. Natural products chemistry and biological properties of the *Ageratum* plant. Toxicol Environ Chem 1995;50:213.
- Kapur SK. Ethnomedicinal plants of Kangra valley (Himachal Pradesh). J Econ Tax Bot 1993;17:395-408.
- Ahluwalia KS. Medicinal plants of Kerala-V, Nagarjun 1968;11:363-9.
- Dutta ML, Nath SC. Ethno-medico botany of Deories of Assam, India. Fitoterapia 1998;69:147-54.
- Gangwar AK, Ramakrishnan PS. Ethnobiological notes on some tribes of Arunachal Pradesh, Northeastern, India. Eco Bot 1990;44:94-105.
- Sudhakar A, Chetty KM. Medicinal importance of some angiospermic weeds used by the rural people of Chittoor district of Andhra Pradesh, India. Fitoterapia 1998;69:390-400.
- Upadhyay OP, Kumar K, Tiwari RK. Ethnobotanical study of skin treatment uses of medicinal plants of Bihar. Pharmaceut Biol 1998;36:167-72.
- Banerjee AK, Banerjee I. A Survey of Medicinal Plants in Shevaroy hills. J Econ Tax Bot 1986;8:271-90.
- Jain S, Pand Puri HS. Ethnomedicinal plants of Jaunsar-Bawar hills, Uttar Pradesh. Indian J Ethnopharmacol 1984;12:213-22.
- Jamir NS. Some interesting medicinal plants used by Nasas. J Res Educ Indian Med 1990;9:81-7.
- Kumar V, Jain SK. A contribution to ethnobotany of Surguja district in Madhya Pradesh, India. Ethnobotany 1998;10:89-96.
- Suresh B, Dhanasekaran S, Kumar RV, Balasubramanian S. Ethnopharmacological studies on the medicinal plants of Nilgiris. Indian Drugs 1995;32:340-52.
- Ramachandran VS, Nair NC. Ethnobotanical observations on Irulars of Tamil Nadu (India). J Econ Tax Bot 1981;2:183-90.
- Maheshwari JK, Singh KK, Saha S. Ethnomedicinal uses of plants by the Tharus of Kheri district, UP. Bull Med Ethnobot Res 1980;1:318-37.
- Siddiqui MB, Alam MM, Husain W. Traditional treatment of Skin diseases in Uttar Pradesh, India. Econ Bot 1989;43:480-6.
- Sankaran S, Alagesaboopathi C. Some Medicinal plants used by the tribals of Shevaroy hills, Tamil Nadu. Flora Fauna 1995;1: 137-8.
- Upadhye A, Kumbhojkar MS, Vartak VD. Observations on wild

- plants used in folk medicine in rural areas of the Kolhapur district. *Ancient Sci Life* 1986;6:119-21.
36. Jain P, Sahu TR. An ethnobotanical study of Noradehi sanctuary park of Madhya Pradesh, India: Native plant remedies for scorpion sting and snake bite. *J Econ Tax Bot* 1993;17:315-28.
 37. Neogi B, Prasad MN, Rao RR. Ethnobotany of some weeds of Khasi and Garo hills, Meghalaya, Northeastern India. *Econ Bot* 1989;43:471-9.
 38. Dhasmana H. Medicinal plants of Pauri town (Garhwal) and adjacent forest regions (part I: family compositae). *J Sci Res Plant Med* 1985;6:45-7.
 39. Rajwar GS. Low altitude medicinal plants of south Garhwal (Garhwal Himalaya). *Bull Med Ethnobot Res* 1983;4:14-28.
 40. Siddiqui MB, Husain W. Some aquatic and marshy land medicinal plants from Hardoi district of Uttar Pradesh. *Fitoterapia* 1992;63:245-8.
 41. Singh H. Ethnobiological treatment of Piles by Bhoxas of Uttar Pradesh. *Ancient Sci Life* 1988;8:167-70.
 42. Dagar HS, Dagar JC. Some folk lore medicinal claims on plants of CarNicobar Island. *Bull Med Ethnobot Res* 1996;17:8-17.
 43. Katewa SS, Arora A. Some plants in folk medicines of Udaipur district (Rajasthan), Ethnobotany 1997;9:48-51.
 44. Sahoo AK, Mudgal V. Ethnobotany of South Chotanagpur (Bihar). *Bull Bot Surv India* 1993;35:40-59.
 45. Sharma PC, Murti KS, Bhat AV, Narayanappa D, Kishore P. Medicinal-Iores of Orissa-I-Skin diseases. *Bull Med Ethnobot Res* 1985;6:93-101.
 46. Hemadri K, Rao SS. Folk lore claims of Koraput and Phulbani district of Orrisa state. *Indian Med* 1989;1:11-3.
 47. Gravena S, Coletti A, Yamamoto PT. Influence of green cover with *Ageratum conyzoides* and *Eupatorium pauciflorum* on predatory and phytophagous mits in citrus. *Bul. OILB-SROP* 1993;16:104-14.
 48. Igoli JO, Ogaji OG, Tor-Anyiin TA, Igoli NP. Traditional Medicine Practice Amongst the Igede people of Nigeria Part-II. *Afr J Traditional Compl Alternative Med* 2005;2:134-52.
 49. Wandji J, Bissangou MF, Ouambra JM, Silou T, Abena A, Keita A. Allelochemicals from *Ageratum conyzoides* L. and *Oryza sativa* L. and their effects on related Pathogens. *Fitoterapia* 1996;67:427.
 50. Devdhar PB, Rao CV. Studies in vegetable oils part-II: Composition of the seed oils of *Eclipta alba* (Linn.) and *Ageratum conyzoides*. *Indian J Appl Chem* 1970;33:305.
 51. Ekundayo O, Laasko I, Hiltunen R. Essential Oil of *Ageratum conyzoides*. *Planta Med* 1988;54:55-7.
 52. Rana VS, Blazquez MA. Chemical composition of the Volatile Oil of *Ageratum conyzoides* aerial parts. *Int J Aromather* 2003;13:203-6.
 53. Rao JT, Nigam SS. Reichst. Aromen Koerperpflegem 1973;23:209-12.
 54. Chalchat JC, Garry RP, Menut C, Lamaty G, Malhuret R, Chopineau J. Correlation between chemical composition and antimicrobial activity VI, Activity of some African essential oils. *J Essent Oil Res* 1997;9:67.
 55. Riaz M, Khalid MR, Chaudhary FM, Rashid KM. Essential oil composition of Pakistani *Ageratum conyzoides* L. *J Essen Oil Res* 1995;7:551.
 56. Pham TT, Nguyen VD, Vien DL. Essential oil of fujian *Ageratum conyzoides* L. *Tap Chi Hoa Hoc* 1976;14:29.
 57. Quijano L, Calderson JS, Gomez GF, Rios T. *In vitro* cytotoxicity of flavonoids against MK2 and C6 tumour cells. *Phytochemistry* 1982;21:2965.
 58. Sharma GP, Garg BD, Girgune JB, Jain NK. Chemical Investigation of the essential oil from *Ageratum conyzoides* Linn. *Univ Indore Res J* 1980a;6:6-11.
 59. Katsuri TR, Manithomes TM. Essential oil of *Ageratum conyzoides*-isolation and structure of two new constituents. *Tetrahedron Lett* 1967;27:2573.
 60. Koteppa P, Rao PJ, Subramanyan B, Rastogi JN, Devakumar C. Benzofuran and other constituents of the Oil of *Ageratum conyzoides*. *Phytochemistry* 1998;49:1385-8.
 61. Pari K, Rao PJ, Subrahmanyam B, Rasthogi JN, Devakumar C. Benzofuran and other constituents of essential oil of *Ageratum conyzoides*. *Phytochemistry* 1998;49:1385-7.
 62. Ahmed AA, Abou-Douh AM, Mohamed AE, Hassan ME, Karchesy J. A New Chromene Glucoside from *Ageratum conyzoides*. *Planta Med* 1999;65:171.
 63. Gonzalez AG, Aguiar ZE, Grillo TA, Luis JG, Rivera A, Calle J. Methoxy Flavone from *Ageratum conyzoides*. *Phytochemistry* 1991;30:1269.
 64. Desai HK, Gawad DH, Govindachari TR, Joshi BS, Kamat VN, Modi JD, *et al*. Chemical Investigation of Indian plants part-VII. *Indian J Chem* 1973;11:840-2.
 65. Srivastava US, Jaiswal AK, Abidi R. Juvenoid activity in extracts of certain plants. *Curr Sci* 1985;54:576-8.
 66. Adesogan EK, Okunade AL. A new flavone from *Ageratum conyzoides*. *Phytochemistry* 1979;18:1863-4.
 67. Adesogan EK, Okunade AL. *J Chem Soc Chem Commun* 1978;152.
 68. Gill S, Mionskowski H, Janczewska D, Kapsa G. Flavonoid compounds of *Ageratum conyzoides* L. herb. *Acta Pol Pharm* 1978;35:241.
 69. Gonzalez AG, Aguiar ZE, Grillo TA, Luis JG, Rivera A, Calle JJ. *Ageratum conyzoides*: A tropical source of medicinal and agricultural products. *Perkin Trans* 1984;1:2945.
 70. Horie T, Tominaga H, Kawamura Y. Revised structure of a natural flavone from *Ageratum conyzoides*. *Phytochemistry* 1993;32:1076-7.
 71. Quijano L, Calderson JS, Gomez GF, Soria IE, Rios T. Highly oxygenated flavonoids from *Ageratum conyzoides*. *Phytochemistry* 1980;19:2439.
 72. Vyas AV, Mulchandani NB. Polyoxygenated flavones from *Ageratum conyzoides*. *Phytochemistry* 1986;25:2625-7.
 73. Nair AG, Kotiyal JP, Subramaian SS. Chemical constituents of the leaves of *Ageratum conyzoides*. *Indian J Pharma* 1977;39:108.
 74. Yadava RN, Kumar S. A novel Isoflavone from the stems of *Ageratum conyzoides*. *Fitoterapia* 1999;70:475-7.
 75. Dubey S, Gupta KC, Matsumoto T, Sterols of *Ageratum conyzoides* L. *Herba Hung* 1989;28:71.
 76. Horng CJ, Lin SR, Chen AH. Phytochemical Study on *Ageratum conyzoides*. *Formosan Sci* 1976;30:101-5.
 77. Hui WH, Lee WK. Triterpenoid and steroid constituents of some *Lactuca* and *Ageratum* species of Hong Kong. *Phytochemistry* 1971;10:899-901.
 78. Wiedenfeld H, Roder E. Pyrrolizidine Alkaloids from *Ageratum conyzoides*. *Planta Med* 1991;57:578.
 79. Sur N, Poi R, Bhattacharyya A, Adityachoudhury NJ. Isolation of aurantiamide acetate from *Ageratum conyzoides*. *Indian J Chem Soc* 1997;74:249.
 80. Pari K, Subrahmanyam B, Rastogi JN, Devakumar C, Rao PJ. Insecticidal (Z)-6-methyl-12-heptadecenoic acid from the essential oil of *Ageratum conyzoides*. *Indian J Chem Sect B* 2000;39B:451.
 81. Riaz M, Khalid MR, Chaudhary FM. *Pak J Sci Ind Res* 1991;34:399.
 82. Tyagi S, Ojha AC, Rawat GS. Chemical examination of some medicinal plants of Shiwalik hills. *Himalayan Chem Pharm Bull* 1994a;11:4-6.
 83. Mondal AK, Parui S, Mandal S. Analysis of the Amino acid content in Pollen of nine Asteraceae species of known Allergenic activity. *Ann Agric Environ Med* 1998;5:17-20.
 84. Tyagi S, Sarraf S, Ojha AC, Rawat GS. Chemical investigation

- of some medicinal plants of Shiwalik hills. *Asian J Chem* 1995;7:165-7.
85. Tyagi S, Saraf S, Ojha AC, Rawat GS. Chemical examination of some medicinal plants (Shiwalik hills). *Himalayan Chem Pharm Bull* 1994b;11:18-21.
 86. Achola KJ, Munenge RW. Activity of *Ageratum conyzoides* on isolated rat phrenic nerve diaphragm and blood pressure on anaesthetised rats. *Int J Pharmacogenetics* 1997;35:31-7.
 87. Marques N, Costalat LT, Fernandes SR, De Napoli MD, Samara AM. Efeito do *Ageratum conyzoides* Lin. Notratamento da artrose. *Rev Bras Rhemaol* 1988;28:109-14.
 88. Whittle SR, Turner A. *J Biochem Pharmacol* 1981;30:1191.
 89. Akah PA. Haemostatic activity of aqueous leaf extract of *Ageratum conyzoides* L. *Int J Crude Drug Res* 1988;26:97.
 90. Agrawal VK. Pharmacognostical studies on the root of *Ageratum conyzoides* L. *J Res Ayur Siddha* 1981;2:242-7.
 91. Gupta A, Gupta R. A survey of plants for presence of cholinesterase activity. *Phytochemistry* 1997;46:827-31.
 92. Achola KJ, Munenga RW, Mwaura AM. Pharmacological properties of root and aerial parts extracts of *Ageratum conyzoides* on isolated ileum and heart. *Fitoterapia* 1994;65:322-5.
 93. Garcia EA, Carvalho MP. Electrophysiological effects of *Ageratum conyzoides* L. on guinea-pig heart. *Phytother Res* 1999;13:172-4.
 94. Marques-Neto JF, Lapa A, Kubota M. Efeitos do *Ageratum conyzoides* Linee no tratamento da artrose. *Rev Bras Reumat* 1988;28:34-7.
 95. Abena AA, Kintangoula-Myaba GS, Diantama J, Bioka D. Effects analgesiques d'un extrait brud' *Ageratum conyzoides* chez le rat. *Encephale* 1993;19:329-32.
 96. Sampson JH, Phillipson JD, Bowery NG, O'Neill MJ, Houston JG, Lewis JA. Ethnomedicinally selected plants as sources of potential analgesic compounds: Indication of *in vitro* biological activity in receptor binding assays. *Phytother Res* 2000;14:24-9.
 97. Okemy-Andissa N, Ouamba JM, Koudou J, Diatewa M, Gleassor H, Abena AA. Comparative Study of Analgesic Activities of Tetra and an Association of Three Plants: *Ageratum conyzoides*, *Combopogon citrates* and *Lippia multiflora*. *Int J Pharmacol* 2006;2:42.
 98. Magalhaes JF, Viana CF, Aragao AG Jr, Moraes VG, Ribeiro RA, Vale MR. Analgesic and anti-inflammatory activities of *Ageratum conyzoides* in rats. *Phytother Res* 1997;11:183.
 99. Silva MJ, Vale MR, Resumos VI. Reuniao Annual da Federac, ao das Sociedades de Biologia Experimental, Caxambu; 1991. p. 296.
 100. Margort deSilva MJ, Capaz FR, Vale MR. Effects of water soluble fraction from leaves of *Ageratum conyzoides* on smooth muscle. *Phytother Res* 2000;14:130.
 101. Mahmood AA, Sidik I, Salmah I, Suzainor KA, Philip K. Antiulcerogenic activity of *Ageratum conyzoides* leaf extract against ethanol-induced gastric ulcer in rats as Animal model. *Int J Mol Med Adv Sci* 2005;1:402-5.
 102. Tandon SK, Chandra S, Tripathi HC. Pharmacological effects of *Ageratum conyzoides* roots, *Indian J Pharma Sci* 1994;56:182.
 103. Moura AC, Silva EL, Fraga MC, Wanderley AG, Afiatpour P, Maia MB. Anti- inflammatory and chronic toxicity study of the leaves of *Ageratum conyzoides* L. in rats. *Phytomedicine* 2005;12:138-42.
 104. Shirwailkar A, Bhilegaonkar PM, Malini S, Kumar JS. The gastroprotective activity of the ethanol extract of *Ageratum conyzoides*. *J Ethnopharmacol* 2003;86:117-21.
 105. Almagboul AZ, Farrog AA, Tyagi BR. Antimicrobial activity of certain Sudanese plants used in folkoric medicine: Screening for antibacterial activity, Part-2. *Fitoterapia* 1985;56:103.
 106. Chah KF, Eze CA, Emuelosi CE, Esimone CO. Antibacterial and wound healing properties of methanolic extracts of some Nigerian medicinal plants. *J Ethnopharmacol* 2006;8:104.
 107. Kishore N, Dubey NK, Tripathi RD, Singh SK. Fungitoxic activity of leaves of some higher plants. *Natl Acad Sci Lett* 1982;5:9-10.
 108. Mishra DN, Dixit V, Mishra AK. Mycotoxic evaluation of some higher plants against ringworm causing fungi. *Indian Drugs* 1991;28:300-3.
 109. Ganesan T, Krishnaraju J. Antifungal properties of wild plants-II. *Adv Plant Sci* 1995;8:194-6.
 110. Mustafa MR, Mahmood AA, Sidik K, Noor SM. Evaluation of wound healing potential of *Ageratum conyzoides* leaf extract in combination with honey in rats as animal model. *Int J Mole Med Adv Sci* 2005;1:406-10.
 111. Oladeio OW, Imosemi IO, Osuagwu FC, Oyedele OO, Oluwadara OO, Ekpo OE, *et al.* A comparative study of the wound healing properties of honey and *Ageratum conyzoides*. *Afr J Med Med Sci* 2003;32:193-6.
 112. Moody JO, Adebisi OA, Adeniyi BA. Do Aloe vera and *Ageratum conyzoides* enhance the anti-microbial activity of traditmedicinal soft soaps (Osedudu)? *J Ethnopharmacol* 2004;92:57-60.
 113. Pandey DK, Asthana A, Tripathi NN, Dixit SN. Volatile plant products vis-à-vis potato pathogenic bacteria. *Indian Perfum* 1981;25:10.
 114. Sharma GP, Jain NK, Garg BD. Antifungal activity of some essential oils-I. *Indian Drugs* 1978;16:21.
 115. Akinyemi KO, Oladapo O, Okwara CE, Ibe CC, Fasure KA. Screening of crude extracts of six medicinal plants used in south-west Nigerian unorthodox medicine for antimethicilin resistant *Staphylococcus aureus* activity. *BMC Compl Alter Med* 2005;5:6.
 116. Jagetia GC, Shirwailkar A, Rao SK, Bhilegaonkar PM. Evaluation of the radioprotective effect of *Ageratum conyzoides* Linn. Extract in mice exposed to different doses of gamma radiation. *J Pharm Pharmacol* 2003;55:1151.
 117. Lans C. Ethnomedicines used in Trinidad and Tobago for reproductive problems. *J Ethnobiol Ethnomed* 2007;3:13.
 118. Abena AA, Ouamba JM, Keita A. Anti-inflammatory, analgesic and antipyretic activities of essential oil of *Ageratum conyzoides*. *Phytother Res* 1996;10:S164-5.
 119. Sharma GP, Jain NK, Garg BD. Antibacterial activity of some essential oils. *Indian Drugs* 1979a;17:14-5.
 120. Sharma GP, Jain NK, Garg BD. Antibacterial activity of some essential oils. *Sci Cult* 1979b;45:327-8.
 121. Pattnaik S, Subramanyam VR, Kole C. Microbioscience 1996;86:237.
 122. Rao JT, Riechst, Aromen, Koerperpflagem 1976;26:50.
 123. Fiori AC, Schwan-Estrada KR, Stangarlin JR, Vida JB, Scapim CA, Cruz ME, *et al.* Antifungal activity of leaf extracts and essential oils of some medicinal plants against *Didymella bryoniae*. *J Phytopathol* 2000;148:483-7.
 124. Maini PM, Morallo-Rejesus B. Toxicity of some volatile oils against golden snail (Pomacea spp.). *Philippines J Sci* 1992;121:391-7.
 125. Asthana A, Chandra H, Dikshit A, Dixit SN. Volatile fungitoxicants from leaves of some higher plants against *Helminthosporium oryzae*. *Pflanzenkr Pflanzenschutz* 1982;89:475.
 126. Singh PP, Shukla HS, Singh RS, Tripathi SC. Antifungal properties of essential oil of *Ageratum conyzoides* L. *Nat Acad Sci Lett* 1986;9:97.
 127. Sharma GP, Jain NK, Garg BD. Antihelmintic activity of some essential oils. *Indian Perfum* 1979c;23:210-2.
 128. Vijayalakshmi K, Mishra SD, Prasad SK. Nematicidal properties of some indigenous plant materials against second stage juvenile of *Meloidogyne incognita* chitwood. *Indian J Entomol* 1979;41:326-31.
 129. Nisar S, Husain SI. Allelopathic effects of some plants on the larval emergence of *Meloidogyne incognita* chitwood. *Indian J Appl Pure*

- Biol 1990;5:129-30.
130. Miller JA, Wood HC. US Patent 1968;121:307.
131. Ragasa CY, Tepora M, Rideout JA. ACGC Chem Res Commun 1998;7:48.
132. Garcia MD, Saenz MT, Gomez MA, Fernandez MA. Topical anti-inflammatory activity of phytosterols isolated from *Erygium foetidum* on chronic and acute inflammation models. *Phytother Res* 1999;13:78-80.
133. Kawaii S, Tomono Y, Katase E, Ogawa K, Yano M. *J Agric Food Chem* 1999;47:128-35.
134. Gbolade AA, Onayade OA, Ayinde BA. Insecticidal activity of *Ageratum conyzoides* L. volatile oil against *Callosobruchus chinensis* F in seed treatment and fumigation laboratory tests: *Insect Science and its application*. *Insect Sci Its Appl* 1999;19:237.
135. Calle J, Rivera A, Luis JG, Aguiar Z, Nimeyer HM, Nathan PJ. Insecticidal activity of the petroleum ether extract of *Ageratum conyzoides* L. *Rev Colomb Quim* 1990;19:91-6.
136. Raja SS, Singh A, Rao SJ. *Anim Morphol Physiol* 1987;34:35.
137. Fagoonee I, Umrit G. Antigonadotropic Hormones from the Goatweed, *Ageratum conyzoides*. *Insect Sci Appl* 1981;4:373-6.
138. Degheele D, Fontier H, Auda M, DeLoof A. *Rijksuniv Gent* 1986;51:101.
139. Halpin RA, Vyas KP, El-Naggar F, Jerina DM. Cytoprotective effects of Modulators of Oxidative Xenobiotic monooxygenase inhibitors. *Chem Biol Interact* 1984;48:297-315.
140. Hammond AH, Garle MJ, Fry JR. *J Biochem Toxicol* 1995;10:265.
141. Ravidranath V, Boyd MR, Jerina DM. *Biochem Pharmacol* 1987;36:441.
142. Darvas B, Jaszberenyi JC, Timar T, Fonagy A. Toxic activity of 2,2-dimethyl-3,4-dichloro- and 2,2-dimethyl-[1,2,3]selenadiazolochromenes on *Pieris brassicae* and larvae *Leptinotarsa decemlineata*. *Pesticide Sci* 1993;18:277-80.
143. Casas J, Gorchs G, Sanchez-Baeza F, Teixidor P, Messeguer A. Inhibition of rat liver microsomal lipid peroxidation elicited by simple 2,2-dimethylchromenes and chromans structurally related to Precocenes. *J Agric Food Chem* 1992;40:585-90.
144. Moreira MD, Picanco MC, Barbosa LC, Guedes RN, Barros EC, et al. Compounds from *Ageratum conyzoides*: Isolation, structural elucidation and insecticidal activity. *Planta Med* 1999;65:2.
145. Atal CK, Srivastava JB, Wali BK, Chakravarty RB, Dhawan BN, Rastogi RP. Screening of Indian plants for biological activity, part-VIII. *Indian J Exp Biol* 1978;16:330-49.
146. Saxena A, Saxena RC. Effects of *Ageratum conyzoides* extract on the development stages of malaria vector, *Anopheles stephensi* (Diptera: Culicidae). *J Env Biol* 1992;13:207-9.
147. Sujatha CH, Vasuki V, Mariappan T, Kalyanasundaram M, Das PK. *Int Pest Control* 1988;30:122-4.
148. Saxena RC, Jayashree S, Padma S, Dixit OP. Evaluation of growth disrupting activity of *Ageratum conyzoides* crude extract on *Culex quinquefasciatus*. *J Env Biol* 1994;15:67-74.
149. Pandey ND, Mathur KK, Pandey S, Tripathi. Effect of some plant extracts against pulse beetle *Callosobruchus chinensis* Linnaeus. *Indian J Entomol* 1986;48:85-90.
150. Bhathal SS, Singh D, Dhillon RS. Insecticidal activity of *Ageratum conyzoides* Linn, against *Lipaphis erysimi* (kaltenbach). *J Insect Sci* 1994;7:35-6.
151. Xu T, Kong C, Hu F. Allelopathy of *Ageratum conyzoides* III: Allelopathic effects of volatile oil from *Ageratum* on plants under different nutrient levels. *Yingyong Shengtai Xuebao* 1999;10:748-50.
152. Hu F, Kong C. Allelopathy of *Ageratum conyzoides* VI, Effects of meteorological conditions on allelopathy of *Ageratum conyzoides*. *Yingyong Shengtai Xuebao* 2002;13:76-80.
153. Kong C, Hu F, Xu X. Allelopathic potential and chemical constituents of volatiles from *Ageratum conyzoides* under stress. *J Chem Ecol* 2002;28:1173-82.
154. Singh AB. Aerobiological studies in India in relation to allergy. *Indian J Pediatr* 1984;51:345-8.
155. Xuan TD, Shinkichi T, Hong NH, Kanh TD, Min CI. Assessment of Phytotoxic action of *Ageratum conyzoides* L. (billy goat weed) on weeds. *Crop Protect* 2004;23:915-22.
156. Arbat A, Patil GV. Common allergenic pollen and spores causing respiratory allergy from central India, Nagpur. *J Soc Pure Appl Nat Sci* 1985;1:5-7.
157. Batish DR, Kohli RK, Singh HP, Saxena DB. Studies on Herbicidal activity of parthenin: A constituent of parthenium hysterothorus towards billy goat weed. *Curr Sci* 1997;73:369-71.
158. Kumar S, Singh CM. Proceedings, Seminar on Control of Lantana and *Ageratum* species. Palampur, India: Himachal Pradesh Agricultural University; 1988. p. 74.
159. Sujatha CH, Nisar S, Jadhi C. Evaluation of plant extracts for biological activity against mosquitoes. *Int Pest Control* 1988;7:122.
160. Kong C, Fei F, Wenju L, Peng W, Yong J. Allelopathic potential of *Ageratum conyzoides* at various growth stages in different habitats. *Allelopathy Jr* 2004;13:233-40.
161. Perumal Samy R, Igancimuthu S, Patric RD. Preliminary Screening of ethnomedicinal plants from India. *J Ethnopharmacol* 1999;66:235-40.
162. Sharma GP, Jain KK, Garg BD. Effect of some essential oils on isolated rectus abdominis of frog (*Rana tigrina*). *Indian Drugs Pharmaceut Ind* 1980b;15:21-2.
163. Vyas AV, Mulchandani NB. Biosynthesis of procoenes-I and II antijuvenile hormones. *Phytochemistry* 1980;19:2597-8.
164. Vyas AV, Mulchandani NB. Structure reinvestigation of conyzorigun: A new chromone from *Ageratum conyzoides*. *J Chem Soc* 1984;2945-7.

Source of Support: Nil, **Conflict of Interest:** None declared.